



# SPF



# INDIANA

June 2012, 12H36

For questions and additional information, please contact:

**Julia Olsen**  
DMHA Bureau Chief  
Mental Health Promotion  
and Addiction Prevention  
(317) 232-7894  
Julia.Olsen@fssa.in.gov

*Prepared for:*  
Indiana Family and Social Services  
Administration  
Division of Mental Health and  
Addiction

*Prepared by:*  
The Center for Health Policy  
Department of Public Health  
Indiana University School of Medicine  
Indiana University-Purdue  
University Indianapolis  
714 N. Senate Avenue, Ste. 220  
Indianapolis, IN 46202



**INDIANA UNIVERSITY**

CENTER FOR HEALTH POLICY  
Department of Public Health

## Indiana Strategic Substance Abuse Prevention Target for 2017

In 2005, the State Epidemiology and Outcomes Workgroup (SEOW) was established as part of the Center for Substance Abuse Prevention's Strategic Prevention Framework State Incentive Grant (SPF SIG) initiative to collect and analyze epidemiological data and facilitate data-based decision-making regarding substance abuse prevention across Indiana. Though the grant funding has ended, the Division of Mental Health and Addiction continues to support the work of the SEOW.

As of this date, the Indiana SEOW has published six annual comprehensive state epidemiological profiles on substance use. *The complete series of The Consumption and Consequences of Alcohol, Tobacco, and Drugs in Indiana: A State Epidemiological Profile* as well as additional documents are available at the Center for Health Policy website at [www.healthpolicy.iupui.edu/SPFSIG/epi](http://www.healthpolicy.iupui.edu/SPFSIG/epi).

Based on the 2011 State Epidemiological Profile, the following substance abuse prevention priorities were identified:

## OUR VISION

Healthy, safe, and drug-free environments  
that nurture and assist all Indiana citizens to thrive.

## OUR MISSION

To reduce substance use and abuse  
across the lifespan of Indiana citizens.

ALCOHOL	TOBACCO	PRESCRIPTION (Rx) DRUGS	MARIJUANA
<p>Reduce past-month alcohol use among 12- to 20-year-olds from 23.6% to 21.2%</p> <p>Data source: NSDUH</p> <p>Target setting method: 10% improvement (Healthy People 2020, objective SA-13.1)</p>	<p>Reduce past-month smoking among adults from 21.2% to 18.0%</p> <p>Data source: BRFSS</p> <p>Target setting method: ITPC, Strategic Plan 2015 (p. 32)</p>	<p>Reduce past-month nonmedical use of prescription pain relievers among high school seniors from 6.6% to 5.9%</p> <p>Data source: ATOD</p> <p>Target setting method: 10% improvement (SEOW consent)</p>	<p>Reduce past-month marijuana use among 18- to 25-year-olds from 16.5% to 14.8%</p> <p>Data source: NSDUH</p> <p>Target setting method: 10% improvement (SEOW consent)</p>
<p>Reduce past-month binge drinking among 12- to 20-year-olds from 17.0% to 15.3%</p> <p>Data source: NSDUH</p> <p>Target setting method: 10% improvement (modified from Healthy People 2020, objective SA-14.1)</p>	<p>Reduce smoking among pregnant women from 18.5% to 12.0%</p> <p>Data source: Indiana Maternal and Child Health Outcomes and Performance Measures Data Book</p> <p>Target setting method: ITPC, Strategic Plan 2015 (p. 32)</p>	<p>Reduce past-month nonmedical use of Rx drugs (excludes pain relievers) among high school seniors from 5.9% to 5.3%</p> <p>Data source: ATOD</p> <p>Target setting method: 10% improvement (SEOW consent)</p>	<p>Reduce past-month marijuana use among 12th grade students from 19.8% to 17.8%</p> <p>Data source: ATOD</p> <p>Target setting method: 10% improvement (SEOW consent)</p>
<p>Reduce past-month binge drinking among 18- to 25-year-olds from 40.8% to 36.7%</p> <p>Data source: NSDUH</p> <p>Target setting method: 10% improvement (modified from Healthy People 2020, objective SA-14.2)</p>			